

## ***Penicillium expansum* toxins correlate with post-harvest fruit rot of apple (*Pyrus malus* L.)**

**Baviskar RN and Dekate HM**

ICLES' Motilal Jhunjunwala College, Vashi, Navi Mumbai, MS, India

### **Manuscript details:**

Received: 31.01.2016  
Accepted: 17.03.2016  
Published : 11.04.2016

**Editor: Dr. Arvind Chavhan**

#### **Cite this article as:**

Dekate HM and Baviskar RN (2016) *Penicillium expansum* toxins correlate with post-harvest fruit rot of apple (*Pyrus malus* L.). *International J. of Life Sciences*, 4(1): 148-150.

**Copyright:** © 2016 | Author(s), This is an open access article under the terms of the Creative Commons Attribution-Non-Commercial - No Derivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

### **ABSTRACT**

Two *Penicillium expansum* toxins viz. patulin and citrinin were found to be associated with *P. expansum* rotted apples. Both toxins were detected in very high amounts. Patulin was detected up to 35 mg/kg whereas citrinin level was comparatively lesser up to 6 mg/kg of the fruit. These amounts are very high in comparison to WHO recommended tolerance level. Present research work, therefore, suggested that proper storage and sorting of apples before marketing is very necessary, so that *P. expansum* rotted fruits are not consumed.

**Key words:** Apples, Mycotoxin, Citrinin, Patulin, Post-harvest.

### **INTRODUCTION**

Apple is an important fresh rosaceous fruit grown in the temperate areas of Jammu and Kashmir. It is juicy, astringent, nutritionally rich in sugars, organic acids, vitamins, Phosphorous, iron and is valued mainly as a dessert fruits. Before harvesting, they are attacked by a few fungi (Chohan and Kaur, 1976) but after harvesting their susceptibility to fungal pathogens increases (Sumbali and Badyal, 1990). During a survey conducted at Vashi fruit market, apples were found to suffer extensive post-harvest blue mold rot caused by *Penicillium expansum*. In view of this, an attempt was made to assess the natural occurrence of *Penicillium* toxins associated with this fruit during pathogenesis.

### **MATERIALS AND METHODS**

#### **Isolation of *P. expansum* isolates from rotted apples:**

Apple suffering from blue mold rot was collected in sterilized bags at regular intervals from wholesaler and retailer. Isolation was made within 24 hours of their collection by streaking the spore on sterilized potato dextrose ager medium.

#### **Preparation of spore suspension:**

Spore suspension of *P. expansum* isolates was prepared by flooding 4 days old culture grown on slant with 10 ml of sterile distilled water containing 0.1% Tween 20. Spore were rubbed from surface of the slant with a glass rod, passed through two layers of sterilized cheesecloth,

counted with a haemocytometer and adjusted with sterile distilled water to obtain  $10^5$  spores/ml.

#### **Inoculation of fruits:**

Mature and healthy apple fruits were initially sterilized by dipping in 95 % ethanol for 30 min., rinsed with sterilized water and then dried under sterilized conditions. Therefore, a single wound 4mm wide and 10mm deep was made in the fruit with the help of a sterilized cork borer and 50  $\mu$ l of spore suspension was inserted into the wound through a sterilized micropipette. Inoculated fruits were incubated at  $28 \pm 2$  °C and 100% relative humidity for 15 days.

#### **Extraction and estimation of patulin and citrinin in from diseased fruit:**

Natural production of patulin and citrinin toxin was estimated at the end of incubation period by taking 30gm of fruit pulps adjacent to the inoculated region and extracting it by the method of Gimeno and Martins (1983). Chloroform extract thus obtained were collected and evaporated to 0.5 ml. This concentrated extract was dissolved to make total volume of 1ml for thin layer chromatography. Quantitative separation of these toxins performed on activated TLC plates using toluene: ethylacetate: chloroform (80:70:50) and 1ml of 90 % formic acid. For detection of patulin, developed plates were sprayed with freshly prepared phenylhydrazine hydrochloride and the heated for 5 minutes at 110°C (Subramanian, 1982). Under visible light, patulin showed yellow spots. For the detection of citrinin, developed plates were directly visualized under 365nm UV light and they showed yellow fluorescent spots. Standard patulin and citrinin samples were also spotted on the TLC plates as reference spots. Further chemical confirmation of citrinin was done by spraying aluminium chloride on the developed plate. Heated it for 5 minutes at 110°C, cooled and observed under 365nm UV light. This chemical test changed the yellow fluorescent of citrinin to sky blue fluorescent (Gimeno and Martins, 1983).

For quantitative estimation of patulin and citrinin, spectrophotometric method of (Bacha *et al.*, 1988) was followed. For patulin estimation, silica gel of each yellow spot was scraped in a clean dry centrifuge tube, dissolved in 5 ml of n-butanol and centrifuged at 3000rpm for 3 min. Citrinin estimation was done by marking yellow fluorescent spots under long wave UV

light (365 nm) scraped them individually in clean, dry centrifuge tubes, dissolved them in 5ml of cold methanol and centrifuged at 3000rpm for 3 minutes. Colour intensity of supernatant was immediately determined in a spectrophotometer at 540nm for patulin and at 332 nm for citrinin.

#### **RESULTS AND DISCUSSION**

Results are depicted in **Table 1** showed that 86.7 % of the recovered isolates were found to be toxigenic producing either patulin or citrinin or both. Among these, positive 33.3 % were found to be patulin producers (28 to 35 mg/kg) whereas, 80.0 % of the tested *P. expansum* isolates were found to be positive for citrinin production (0.09 to 6.04 mg/kg). Similar results were obtained by (Brian *et al.*, 1956). The production of patulin in fruits may be attributed to the sugar, organic acid (Menniti *et al.*, 2010).

From the result it was also observed that 26.7 per cent isolates of *P. expansum* could produce both patulin and citrinin and their levels of production were very high. Similarly, level of patulin and citrinin contamination from *P. expansum* infected apples has been reported to be as high as 1 and 3 mg/kg (Gimeno and Martins, 1993; Ritieni, 2003). Although there is currently no evidence to prove that patulin and citrinin have the potential to produce adverse human health effects (Hopkins, 1993; Machinosky and Midio, 1995), yet the finding in animals that these mycotoxins are carcinogenic, teratogenic and nephrotoxic (Krough *et al.*, 1970; Dickens and Jones, 1961; Mayer and Legator, 1969), emphasize the need for concern. World health Organisation (WHO) has recommended a maximum patulin level of 50 $\mu$ g/kg. Various European and other countries have also recommended maximum patulin levels of 30-50  $\mu$ g/kg (Italian Ministry of Health, 1999). But so far no recommendation for the tolerance level of citrinin has been made. Since patulin is a stable mycotoxin (Poland and Allen, 1970; Harrison, 1989), and its co-occurrence with citrinin in the infected crab apples and the products made from crab apples rotted by *P. expansum*, which may present a health hazard.

**Table 1: Production of patulin and citrinin in *P. expansum* infected apples**

Total no. of isolates screened= 15.0		
	% of toxigenic isolates	= 86.7
Isolates screened	% of patulin producer	= 33.3
	% of citrinin producer	= 80.0
	% of both patulin and citrinin	
	Producer	= 26.7
	Patulin toxin	Citrinin toxin
Pe-1	-	6.04 ± 7.1
Pe-2	29.50±6.5*	0.94 ± 2.0
Pe-3	-	0.26 ± 1.7
Pe-4	-	0.09 ± 2.4
Pe-5	-	0.72 ± 1.8
Pe-6	-	0.15 ± 3.4
Pe-7	-	4.21 ± 2.4
Pe-8	-	0.39 ± 1.8
Pe-9	-	-
Pe-10	28.00±3.5	1.15 ± 6.4
Pe-11	33.50±7.5	-
Pe-12	-	-
Pe-13	35.00±5.8	0.14 ± 2.4
Pe-14	-	0.18 ± 1.4
Pe-15	33.50±8.5	2.00 ± 1.6

\*Each value represents the mean ± SD, n = 3, and - not detected.

This evaluation, therefore, shows that proper storage and sorting of apples is very necessary before marketing so that intake of these mycotoxins above the tolerance limit is avoided. This will essentially make a difference for children and vegetarians who consume a higher quantity of fruits.

## REFERENCES

- Bacha HR, Hadidane EE, Creppy C, Regnault FE and Dircheimer G. (1988) Monitoring and identification of fungal toxins in food products, animal feed and cereals in Tunisia. *Jour. Stored Prod. Res.* 24(4): 199-206.
- Brian PW, Elson GW and Lowe D (1956) Production of patulin in apple fruits by *Penicillium expansum*. *Nature*, 178: 263-264.
- Chohan JS and Kaur S (1976) Occurrence of Indian crab apple scab in plains of Punjab. *Indian Phytopathol.* 29: 441-442.
- Dickens F and Jones HE (1961) Carcinogenic activity of a series of reactive lactones and related substances. *Br. Jour. Cancer.* 15: 85-100.
- Gimeno A and Martins ML (1983) Rapid thin layer chromatography determination of patulin, citrinin, and aflatoxin in apples and pears and their juices and jams. *Jour. Assoc. Off. Anal. Chem.* 66(1): 85-91.
- Harrison MA (1989) Presence and stability of patulin in apple products: a review. *Jour. Food Safety.* 9: 147- 153.
- Hopkins J (1993) The toxicological hazards of patulin. *Food Chem. Toxicol.* 31: 455-456.
- Italian Ministry of Health (1999) Maximal acceptable mycotoxin concentrations in foods of national and international origin. Circular dated June 9<sup>th</sup>.
- Krough P, Hasselager E and Friis P (1970) Isolation of two nephrotoxic compounds from *P. viridcatum* Westling: Citrinin and oxalic acid. *Acta Pathol. Microbiol Scand Sect B.* 78: 401-413.
- Machinsky MJ and Midio AF (1995) Toxicological and analytical aspects of patulin in foods. *Rev. Farm. Biogum. Univ. S. Paulo.* 31: 1-19.
- Mayer VW and Legator M (1969) Production of petite mutant of *Sacchromyces cerevisea* by patulin. *Jour. Agrl. Food Chem.* 17: 454- 456.
- Menniti AM, Neri F, Gregor R and Maccaferri M (2010) Some factors influencing patulin production by *Penicillium expansum* in pome fruits. *Jour. Sci. Food Agri.* 90(3): 2183-2187.
- Poland AE and Allen R (1970) Stability studies with patulin. *Jour. Assoc. Off. Anal. Chem.* 53: 688-691.
- Ritieni A (2003) Patulin in italian commercial apple products. *Jour. Agric. Food Chem.* 51: 6086-6090.
- Subramanian T (1982) Colorimetric determination of patulin produced by *Penicillium patulum*. *Jour. Assoc. Off. Anal. Chem.* 65: 5-7
- Sumbali G and Badyal K (1990) Some new market diseases of pome fruits from India. *Nat. Acad. Sci. Letters.* 13: 331-332.