



P.V. Sukhatme: A Pioneer in the Application of Statistics to Agriculture and Nutrition

Kharade RN¹ and Shinde MB²

¹Department of Statistics, | ²Department of Mathematics, Shahajiraje Mahavidyalaya, Khatav, Shivaji University, Kolhapur. MS, India.

*Corresponding Author : Shinde MB

Email: maheshshinse4833@gmail.com

Manuscript details:

Received: 30.11.2025
Accepted: 28.12.2025
Published: 31.12.2025

Cite this article as:

Kharade RN and Shinde MB (2025)
P.V. Sukhatme: A Pioneer in the
Application of Statistics to Agriculture
and Nutrition, *Int. J. of Life Sciences*, 13
(4): 473-476.

Available online on <http://www.ijlsci.in>
ISSN: 2320-964X (Online)
ISSN: 2320-7817 (Print)



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third-party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

ABSTRACT

A pivotal figure in twentieth-century statistics, Pandurang Vasudeo Sukhatme significantly impacted national systems and international politics through his advancements in human nutrition and agricultural survey methods. The General Crop Estimation Surveys in India and other national programs that have been adopted throughout Asia and Africa were made possible by his leadership at ICAR, which formalized crop-cutting surveys and stratified multi-stage sampling for yield estimation. The lower-cutoff approach in FAO's Prevalence of Undernourishment (PoU) methodology, which continues to be refined today, was influenced by his stochastic model of intra-individual energy requirements, which reframed the measurement of undernourishment and sparked the shift from protein-centric narratives to energy adequacy. This study examines the methodological foundation of Sukhatme's work, its institutional dissemination, scientific criticisms, and policy implications, emphasizing its ongoing significance for nutrition and food security statistics.

Keywords: P. V. Sukhatme; crop-cutting surveys; stratified multi-stage sampling; intra-individual variation; FAO undernourishment; IASRI; agricultural statistics; energy requirements.

INTRODUCTION

Pandurang Vasudeo Sukhatme was born on July 27, 1911, in Budh, Satara, Maharashtra (Narain, 1995). He earned a B.Sc. from Fergusson College in 1932 before pursuing Ph.D. in 1936 and a D.Sc. degree (1939) at University College London, where he worked under R. A. Fisher, Jerzy Neyman, and E. S. Pearson (Narain, 1995; Pandurang Vasudeo Sukhatme - biographical and publishing profile). He then supervised agricultural statistics reforms at the Indian Council of Agricultural Research (ICAR) from 1940 to 1951 and managed the Statistics Division at the Food and Agriculture Organization (FAO) from 1951 to 1971 (Narain, 2000; IASRI Vision 2050). His work combined rigorous statistical theory with practical policy, notably in agricultural output estimation and undernutrition assessment (Sukhatme; Panse, 1970-1974; Sukhatme, 1961).

Contributions to Agricultural Statistics

1. The problem context and design innovation

Pre-reform agricultural statistics in India depended mainly on administrative judgments with no probabilistic basis, resulting in distorted and delayed estimates (Sukhatme, 1945; Sukhatme, 1945 Interim Report). To address these shortcomings, Sukhatme developed and scaled stratified multi-stage random sampling for yield estimation, which is often used in crop-cutting surveys (CCEs) to objectively estimate average yields while measuring sample error (Sukhatme, 1945; Sukhatme; Panse, 1970/1974). His early demonstrations, which included rice yield estimation at the province level, laid the groundwork for randomized plot selection within a probability framework (Sukhatme, 1945).

2. Crop-cutting surveys and national systems

With the support of initiatives like the Timely Reporting Scheme (TRS) and the Establishment of an Agency for Reporting Agricultural Statistics (EARAS), which standardized sample frames, timing, and supervision across states, the methodology developed into a national system through the General Crop Estimation Surveys (GCES) (MoSPI, 2008; IASRI Vision 2050). Unbiased yield calculations at scale were made possible by design features such as sub-district stratification, random village selection from TRS samples, two independent CCEs per chosen village, per crop, and allocation proportionate to farmed area (MoSPI, 2008).

The spread of sample survey methodology in fertilizer quality, fisheries, horticulture, post-harvest loss estimation, and agricultural insurance (small area estimation) is monitored by the Indian Agricultural Statistics Research Institute (IASRI), the institutional successor to the ICAR statistical unit that Sukhatme assisted in establishing (IASRI Vision 2050; Sukhatme; Panse, 1970-1974).

3. Theoretical codification and global influence.

Sukhatme turned experience into theory in *Sampling Theory of Surveys with Applications*, a widely used reference that linked ratio/regression estimators, multistage designs, and nonsampling error concerns to field implementation (Sukhatme; Panse, 1970/1974). IASRI's institutional roadmap attributes these advances to national and worldwide acceptance across sectors and nations, highlighting the long-term influence of the "Sukhatme era" in agricultural statistics (IASRI Vision 2050; Narain, 2000).

Nutrition and the Stochastic Model of Energy Requirements

1. From protein deficiency to energy adequacy

Moving to FAO, Sukhatme reframed human nutrition measurements by prioritizing energy sufficiency over protein-centric narratives common in mid-century discourse (Sukhatme, 1961; Ferro-Luzzi, 1990). His 1961 JRSS work on world hunger examined energy balance at a large scale and anticipated policy disputes on how to define and assess malnutrition (Sukhatme, 1961). The latter FAO method was built on the pivot toward dietary energy consumption rather than a set, universal need rate (FAO, 2009; FAO, 2016).

2. Intra-individual variation and homeostatic regulation

Sukhatme's nutrition contribution is based on the hypothesis that individual dietary energy requirement (DER) is a stochastic, stationary process with significant intra-individual variation and autoregulatory homeostasis, allowing for short-run intake fluctuations without long-term weight change (Sukhatme; Margen, 1982; AJCN, 1985). The seminal AJCN study with Margen proposed that the coefficient of variation in intake can be significant with body weight stability, meaning that "requirement" is dynamic and energy balance varies around zero within a homeostatic range (Sukhatme; Margen, 1982). This study provides statistical support for employing a lower threshold (representing the lower bound of the requirement distribution) rather than a population mean requirement to detect chronic malnutrition (Sukhatme, 1961; Sukhatme; Margen, 1982; FAO, 2009).

3. FAO's methodology and subsequent refinements

The FAO transformed this insight into a univariate framework for the Prevalence of Undernourishment (PoU), estimating the fraction of the population whose usual dietary energy consumption falls below a lower requirement cutoff derived from demographic and physical activity characteristics, body weight distributions, and basal metabolic rates. FAO continues to refine distributional assumptions, cutoff derivations, and data inputs while maintaining the critical intuition that some observed consumption variability is due to normal requirement variability rather than food insecurity (FAO Open Knowledge, 2014-2023; Smith, R.; FAO/IFPRI literature).

The FAO's methodological retrospectives explicitly trace this lineage back to Sukhatme's model, while also addressing criticisms by clarifying the role of joint versus marginal distributions and improving component variance estimation for the cutoff (FAO Exploding the Myth of the Bivariate Distribution; Kakwani, 1986; 2015).

Methodological Controversies and Evidence

1. The lower-cutoff debate

Critics have claimed that the univariate lower-cutoff technique may underestimate malnutrition by disregarding the risk inside the overlapping area of intake and requirement distributions or by overemphasizing adaptation (Chakrabarti, 1981; Kakwani, 2015). FAO and subsequent theoretical work demonstrated that the bivariate framework reduces to a univariate cutoff when positive correlation and unimodality assumptions are met, thus justifying the lower-cutoff method for policy statistics while encouraging better specification of requirement variability (FAO Exploding the Myth of the Bivariate Distribution; Kakwani, 1986). The argument honed the empirical agenda for requirement variance components and the intensity of intake-requirement correlation (Kakwani, 2015; Chakrabarti, 1981).

2. Evidence on intra- vs inter-individual variability

Metabolic balance and intake data analysis revealed significant intra-individual variability with limited long-term weight drift in steady environments, which is consistent with autoregulatory mechanisms (AJCN, 1985; Sukhatme; Margen, 1982); these findings supported the stationary requirement hypothesis and motivated autoregressive modeling of day-to-day energy balance (Kakwani, 1986; AJCN, 1985). While prominent nutritionists disputed the extent of intra-individual variance and the public health implications of "adaptation," the statistical case for a lower bound cutoff became stronger as FAO decomposed requirement variance into demographic, body-weight, activity, and residual components and operationalized these within PoU estimation.

Policy and Institutional Impact

1. Agriculture: from subjective assessment to scientifically audited systems

India adoption of GCES, TRS, EARAS, and rigorous CCE procedures re-based national foodgrain estimate on

reproducible, auditable designs, minimizing bias and allowing credible policy responses to supply shocks and price stability requirements (MoSPI, 2008; IASRI Vision 2050). IASRI's adoption of survey methods in sectors such as horticulture, fisheries, and post-harvest loss estimation exemplifies a broad system transformation traceable back to Sukhatme's design philosophy (IASRI Vision 2050; Sukhatme; Panse, 1970/1974). These frameworks have been investigated and implemented in a variety of Asian and African contexts, demonstrating their resilience and cost-effectiveness for official statistics (Narain, 2000; IASRI Vision 2050).

2. Nutrition: the architecture of global hunger measurement

FAO's PoU, used in World Food Surveys, World Food Conference assessments, and State of Food Insecurity in the World reports, is based on Sukhatme's lower-cutoff tradition, with modern refinements in response to data and methodological advances (FAO, 2009; FAO, 2016; FAO Open Knowledge, 2014-2023). The technique continues to underlie SDG hunger monitoring, with FAO advice highlighting the conceptual separation of normal need fluctuation and food insecurity while enhancing country-level parameterization (FAO, 2009; FAO, 2016). The ongoing discussion meant that the approach was evaluated and continuously improved, and also increased statistical literacy in nutrition policy circles (Kakwani, 2015; Smith, R.).

Enduring Legacy and Relevance

Sukhatme combined theory, design, and policy in two areas where measurement error has first-order consequences: food production and malnutrition (Narain, 1995; Sukhatme, Panse, 1970-1974). In agriculture, he connected national statistics architecture with probability sampling methodology, allowing estimation error to be quantified and managed in real time (MOSPI, 2008; IASRI Vision 2050). In nutrition, he changed the paradigm from fixed norms to stochastic requirements, which influenced a measurement system used for decades in worldwide monitoring (Sukhatme, 1961; FAO, 2009). Current talks on small-area estimation for crop insurance, remote-sensing-integrated sample frames, and enhanced PoU parameterizations remain within the conceptual domain he helped define (IASRI Vision 2050; FAO Open Knowledge, 2014,2023).

CONCLUSION

P. V. Sukhatme's methodological clarity and institutional leadership resulted in long-lasting statistical systems in agriculture and nutrition, which are still functioning and evolving. Crop-cutting and GCES architecture altered yield estimation, informing similar studies across sectors and geographies. The FAO's global hunger metrics are based on a stochastic perspective of energy requirements, with the lower-cutoff method supported by theoretical rationale and empirical modifications. His work reveals how rigorous statistical thinking may redirect national programs and international policy towards more accurate and actionable assessment.

Acknowledgments

The authors acknowledge institutional documentation and methodological manuals that preserve and extend Sukhatme's contributions within IASRI, FAO, and India's national statistical system

Conflict of Interest: The authors declare no conflict of interest in relation to this research.

Data Availability Statement: Not applicable.

Correspondence and requests for materials should be addressed to **Shinde MB**

Peer review information

IJLSCI thanks the anonymous reviewers for their contribution to the peer review of this work. A peer review file is available.

Reprints and permissions information is available at <https://www.ijlsci.in/reprints>

REFERENCES

- Narain, P. (1995). Pandurang Vasudeo Sukhatme. Biographical Memoirs of Fellows of the Indian National Science Academy, 20, 147-167.
- Narain, P. (2000). On some contributions of Pandurang Vasudeo Sukhatme. Journal of the Indian Society of Agricultural Statistics, 53(3), 229-246.
- Sukhatme, P. V.; Margen, S. (1982). Autoregulatory homeostatic nature of energy balance. American Journal of Clinical Nutrition, 35(2), 355-365.
- FAO (2009). Methodology for estimating the prevalence of undernourishment (PoU) background notes and exposition.
- Sukhatme, P. V. (1945). Random sampling for estimating rice yield in Madras Province. Indian Journal of Agricultural Science, 15(3), 119-129.
- Sukhatme, P. V. (1961). The worlds hunger and future needs in food supplies. Journal of the Royal Statistical Society, Series A, 124(4), 463-525.
- FAO (Exploding the Myth of the Bivariate Distribution). Review and justification of the univariate-cutoff framework and debate history.
- Sukhatme, P. V. (1945). Interim report on crop-cutting survey for estimating yield.
- Ferro-Luzzi, A. (1990). Social and public health issues in adaptation to low energy intake. American Journal of Clinical Nutrition context and debates.
- FAO (2016). The FAO method to estimate the Prevalence of Undernourishment (technical presentation).
- MoSPI (2008). Manual on Area and Crop Production Statistics (TRS, EARAS, GCES design and procedures).
- AJCN (1985). Long-term metabolic balance studies and implications for energy balance (discussion of Sukhatme Margen).
- Smith, R.; FAO/IFPRI related literature on requirement cutoffs and PAL/BMR assumptions in PoU (book chapter).
- Sukhatme, P. V.; Panse, V. G. (1970/1974). Sampling Theory of Surveys with Applications (Iowa State/ISAS editions).
- Later physiology literature contextualizing energy balance regulation (thermogenesis, adipose tissue roles).
- Chakrabarti, S. K. (1981). Measurement of incidence of undernutrition (critical discussion of models).
- Pandurang Vasudeo Sukhatme biographical and publications summary (career chronology; contributions list).
- FAO Open Knowledge (2014-2023). Refinements to FAO PoU methodology and responses to critiques.
- Statistics Day (India) commemorative booklet (biographical note).
- Kakwani, N. (2015). Measuring food insecurity (critique and theoretical clarification; correlation conditions for univariate reduction).
- Kakwani, N. (1986). On measuring undernutrition (variance decomposition; autoregressive modeling references).
- IASRI Vision 2050: Diffusion of agricultural survey methodologies across sectors and countries.

© The Author(s) 2025

Publisher's Note

IJLSCI remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.