



Unveiling Breast Cancer ChemoPreventive Phytochemicals: GC-MS Profiling and Antioxidant Potentials of *Indigofera longiracemosa* Boivin ex Bail

K. Shabna^{1*} and G. Arun Kumar²

¹Ph. D. Scholar | ²Assistant Professor, Department of Microbiology, Hindusthan College of Arts & Science, Coimbatore, India, 641 028

*Corresponding author email : shabnakalathil@gmail.com¹ | arunkumar.g@hicas.ac.in²

Manuscript details:

Received: 09.03.2026
Accepted: 27.03.2026
Published: 30.03.2026

Cite this article as:

K. Shabna and G. Arun Kumar (2026)
Unveiling Breast Cancer
ChemoPreventive Phytochemicals: GC-
MS Profiling and Antioxidant Potentials of
Indigofera longiracemosa Boivin ex Bail,
Int. J. of Life Sciences, 14 (1): 15-23.

Available online on <http://www.ijlsci.in>
ISSN: 2320-964X (Online)
ISSN: 2320-7817 (Print)



Open Access This article
is licensed under a
Creative Commons

Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third-party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

ABSTRACT

High recovery of various phytochemicals, such as alkaloids, terpenoids, fatty acids, and phenylpropanoids, was confirmed by acetone extraction of 400 g of plant material, yielding 60 g of crude extract (15% w/w). Strong antioxidant activity was demonstrated by the DPPH radical scavenging assay, which showed concentration-dependent absorbance decline (0.431-0.171) and % RSA increase (53.7-80.3%) at 20-100 µg/ml; linear regression ($Y=0.336X+46.89$; $R^2=0.993$) determined $IC_{50}=9.25$ µg/ml, which was better than the BHT standard ($IC_{50}=12$ µg/ml; $Y=0.384X+45.392$; $R^2=0.9999$). Alpha-linolenic acid (9,12,15-octadecatrienoic acid), phytol, neophytadiene, trans-indigo, 1H-indole-2,3-dione, and alpha-tocopherol-β-D-mannoside, many of which have been shown to have antioxidant, anti-inflammatory, cytotoxic, and anticancer effects relevant to breast cancer via apoptosis induction, proliferation inhibition, and oxidative stress modulation. This phytochemical synergy highlights the extract's multi-target potential against oxidative damage and carcinogenesis, positioning it as a promising chemopreventive candidate.

Keywords: Phytochemicals, Antioxidant activity, DPPH Assay, IC_{50} , GC-MS, Chemopreventive.

INTRODUCTION

With an anticipated 19.3 million new cases and 10.0 million cancer-related deaths in 2020, cancer is one of the main causes of death globally (Sung *et al.*, 2021). The unchecked proliferation and spread of aberrant cells are hallmarks of cancer, a complex and multidimensional process (Bray *et al.*, 2018). Genetic changes that impair normal cell division and growth are one of the many causes of cancer (Siegel *et al.*, 2020). While chemotherapy drugs can prolong life expectancy and offer short-term comfort (Weissenstein *et al.*, 2014). Several anticancer drugs have adverse side effects. Traditional medicine has employed plant-derived substances for millennia, and many of them have demonstrated promise

in the treatment of cancer (Pan *et al.*, 2013). Natural substances have long been used to cure and prevent a wide range of illnesses (Singh *et al.*, 2012), serving as the foundation of the conventional healing system (Dehelean *et al.*, 2021). Furthermore, a larger population may be able to obtain cancer therapy more easily thanks to the utilization of plant extracts as a sustainable and affordable substitute for pricey and frequently unavailable pharmaceutical medications (Mukherjee *et al.*, 2018).

Typically, phytochemicals or physiologically active substances with varying polarity combine to form plant extracts. Separation continues to be a significant obstacle in the identification and classification of physiologically active chemicals (Altemimi, 2017). A combination analytical method called gas chromatography-mass spectroscopy (GC-MS) is utilized to identify and ascertain the substances found in a plant sample (Uma & Balasubramaniam, 2012). GC-MS is crucial for phytochemical analysis and chemotaxonomic research on medicinal plants that have components that are physiologically active (Hethelyi *et al.*, 1987).

The Fabaceae family includes the small tree or shrub *Indigofera longiracemosa* Boivin ex Baill, which is indigenous to India, Sri Lanka, and Nepal. It has a cylindrical crown and is between two and five meters tall. It can be found in scrublands and dry deciduous woodlands. In rats, the roots have been shown to have antidiabetic benefits (Kumar *et al.*, 2015), neutralize all snake venoms (Thangadurai, 1998), and have antioxidant (Patel *et al.*, 2013). Leaves have traditionally been used to treat wounds and skin conditions (Sebaugh, 2011). Leaves have antibacterial qualities against certain bacterial strains. The antifungal qualities of seeds can affect some fungus strains (Kumar *et al.*, 2009). However, no studies on *I. longiracemosa* anticancer potential have been conducted. Because of this evidence, *I. longiracemosa* was my choice for this investigation.

This study involved a sequential analysis of the acetonic extract of *Indigofera longiracemosa* Boivin ex Baill (AEIL), starting with qualitative phytochemical screening, followed by evaluation of antioxidant potential using the DPPH radical scavenging assay and concluding with GC-MS analysis to identify specific bioactive compounds.

MATERIALS AND METHOD

Plant Collection and Extraction: The leaves of *I. longiracemosa* were collected in October 2022 from the Edayur north area of the Malappuram district of Kerala (Latitude = 10.88035, Longitude = 76.09092). In Coimbatore, Tamil Nadu, India, Dr M.U. Sharief, Scientist 'F', Botanical Survey of India, Southern Circle, Coimbatore, verified the validity of the plant. After 400 g of plant material was shade-dried, it was removed using 80% acetone and left to stand at room temperature for 72 hours. At a regulated temperature of 40 to 50 °C and at a lower pressure, the extract was filtered and concentrated in a rotating evaporator until it was dry. The 60 g dark-brown solid extract was kept in a vacuum desiccator at 4 °C until needed again.



Fig. 1 *Indigofera longiracemosa* Boivin ex Baill

Preliminary examination of phytochemicals

The extract was examined using common screening techniques to determine whether it contained different phytochemical ingredients (Wagner *et al.*, 1984). For the detection of terpenoids, phenols, steroids, alkaloids, tannins, flavonoids, glycosides, etc., a standard procedure was employed.

Antioxidant activity by DPPH Assay

Procedure: The method suggested by (Blois, 1958) was used to assess the plant extracts' antioxidant activity against DPPH. A DPPH dilution of 1×10^{-4} M was made in methanol. Each sample was separated into aliquots of 1 mL in the methanolic extract (at four distinct concentrations: 0.1, 0.5, 1, and 2 mg/mL; two replicates per sample and concentration), and 2 mL of DPPH methanolic dilution was then added. After 16 minutes at room temperature in the dark, the absorbance of the mixture was measured at 517 nm using a UV spectrophotometer. The blank was made

using DPPH diluted with methanol. The antioxidant activity (% inhibition or % scavenging) is calculated using the formula.

$$\text{DPPH Scavenging Activity (\%)} = \frac{(A_{\text{control}} - A_{\text{sample}})}{A_{\text{control}}} \times 100$$

Standard for DPPH: One of the most widely used antioxidants, butylated hydroxytoluene (BHT), is suitable for use in DPPH. This can be used as a reference standard for comparing the antioxidant activity of plant extracts.

IC₅₀ Calculation : The IC₅₀ value for both sample and standard was calculated using the equation: **IC₅₀ = (50-b)/m**, where b is the intercept and m is the slope of the linear regression line (Sebaugh, 2011).

Gas chromatography- mass spectroscopy (GC-MS) - A Shimadzu Nexis GC 2030 system with an AOC 30/20i autosampler managed by GCMS Solutions software was used for gas chromatography-mass spectrometry (GC-MS) analysis. Samples were automatically added to the system by the autosampler, and separation was accomplished on a 30-meter-long SH I-5Sil MS capillary column with an inner diameter of 0.25 mm and a film thickness of 0.25 µm that was appropriate for low-polarity compounds. To maximize peak resolution while reducing analysis time, the oven

temperature was set in a stepwise ramp. Other inert gases could also be used, but helium was the carrier gas (Hubschmann, 2015).

RESULTS & DISCUSSION

The extraction of 400 g of plant material yielded 60 g of extract, resulting in an extraction efficiency of 15% (w/w). Acetone extraction yielded a diverse range of phytochemicals, including alkaloids, terpenoids, fatty acids, diterpenoids, triterpenoids and other compounds, indicating its effectiveness in extracting a broad spectrum of bioactive compounds.

The DPPH radical scavenging activity of the sample and standard (BHT) was calculated at various concentrations (20-100 µg/ml). The absorbance values of the sample decreased with increasing concentration, with values ranging from 0.431 to 0.171. The percentage RSA (%) of the sample increased with increasing concentration, with values ranging from 53.7 to 80.3. The IC₅₀ values were calculated using linear regression analysis of the concentration- response data. For the sample, an IC₅₀ value of 9.25 µg/ml, indicating its potent antioxidant activity. The results are presented in Table 1 and Figure 2.

Table 1: DPPH Radical Scavenging Activity of the Sample

Sl.no	Concentration(µg/ml)	Absorbance of the sample	% RSA	IC ₅₀
1	20	0.431	53.7	9.25
2	40	0.373	59.6	
3	60	0.326	64.4	
4	80	0.242	73	
5	100	0.171	80.3	






Figure 2: DPPH Radical Scavenging Activity of the sample

Table 2: DPPH Radical Scavenging Activity of the standard (BHT)

Sl.no	Concentration($\mu\text{g/ml}$)	Absorbance of the BHT	% RSA	IC50
1	20	0.485	47.6	12
2	40	0.412	55.5	
3	60	0.342	63.1	
4	80	0.272	70.6	
5	100	0.202	78.2	

**Figure 3: DPPH Radical Scavenging Activity of the standard (BHT)****Figure 4: GCMS Profiling**

In comparison, the standard BHT exhibited absorbance values ranging from 0.485 to 0.202, with % RSA values ranging from 47.6 to 78.2 %. The IC₅₀ value of BHT was 12 $\mu\text{g/ml}$. These results indicate that the sample exhibits antioxidant activity, with a lower IC₅₀ value compared to BHT, suggesting higher antioxidant potency. The high R^2 values indicate a strong correlation between concentration and

response for both the sample and BHT. The results are presented in Table 2 and Figure 3.

The GCMS analysis identified 25 chromatographic peaks in the sample, with retention times, relative peak area percentages, and compound assignments. The data is represented in Figure 4. Those resolved into 17 unique compounds after merging duplicates.

Repeated peaks indicate a high abundance of certain constituents.












The bioactive compounds identified in the GCMS exhibit a range of properties relevant to cancer research, including antioxidant, antimicrobial, and potential anticancer activities. Alpha, -tocopherol-.beta. -D-mannoside, a vitamin E derivative, is a potent

antioxidant that may help protect normal cells from chemotherapy-induced oxidative stress and potentially alleviate oxidative damage in cancer cells. Trans-indigo, a natural indole pigment, exhibits antimicrobial properties and has been investigated for its anti-leukemic and other anticancer effects, inducing apoptosis in certain cancer cell lines.

Table 3: GC-MS Peak Data for Compound Identification

Peak#	R.Time	Area	Area%	Height	Height%	A/H	Mark	Name
1	7.645	98501	1.33	31190	1.42	3.16	MI	Isoamyl ethanoate
2	16.140	155828	2.11	47359	2.15	3.29	MI	Anthranilic acid
3	23.646	869428	11.75	262952	11.95	3.31	MI	1H-Indole-2,3-dione
4	24.787	165425	2.24	61646	2.80	2.68	MI	trans-1,2-Diphenylcyclobutane
5	25.580	60960	0.82	20934	0.95	2.91	MI	2,3-Bis(1-methylallyl)pyrrolidine
6	26.697	283112	3.83	103410	4.70	2.74	MI	Neophytadiene
7	27.120	90704	1.23	34036	1.55	2.66	MI	Anthracene, 9-ethenyl-
8	27.580	90944	1.23	36211	1.65	2.51	MI	Neophytadiene
9	28.189	580386	7.85	161955	7.36	3.58	MI	3-Hydroxy-3-methyl-oxindole, Ac derivative
10	29.145	347916	4.70	130035	5.91	2.68	MI	n-Hexadecanoic acid
11	32.009	128501	1.74	36132	1.64	3.56	MI	Phytol
12	32.459	314014	4.25	99634	4.53	3.15	MI	9,12,15-Octadecatienoic acid, (Z,Z,Z)-
13	36.120	124933	1.69	39723	1.81	3.15	MI	Cyclononasiloxane, octadecamethyl-
14	37.652	901227	12.18	307924	14.00	2.93	MI	Cyclohexane, 1,3,5-triphenyl-
15	38.297	121507	1.64	43976	2.00	2.76	MI	Cyclononasiloxane, octadecamethyl-
16	39.059	403925	5.46	131973	6.00	3.06	MI	(2,3-Diphenylcyclopropyl)methyl phenyl sulfoxide, trans-
17	39.288	1157882	15.65	254761	11.58	4.54	MI	(2,3-Diphenylcyclopropyl)methyl phenyl sulfoxide, trans-
18	39.385	180876	2.45	66360	3.02	2.73	MI	(2,3-Diphenylcyclopropyl)methyl phenyl sulfoxide, trans-
19	39.577	242709	3.28	84435	3.84	2.87	MI	(2,3-Diphenylcyclopropyl)methyl phenyl sulfoxide, trans-
20	40.345	147679	2.00	45203	2.06	3.27	MI	Tetracosamethyl-cyclododecasiloxane
21	41.010	90554	1.22	29522	1.34	3.07	MI	(2,3-Diphenylcyclopropyl)methyl phenyl sulfoxide, trans-
22	42.256	138189	1.87	37918	1.72	3.64	MI	Tetracosamethyl-cyclododecasiloxane
23	44.043	120504	1.63	30389	1.38	3.97	MI	Tetracosamethyl-cyclododecasiloxane
24	45.167	249340	3.37	53279	2.42	4.68	MI	trans-Indigo
25	48.426	331932	4.49	48631	2.21	6.83	MI	. Alpha.-Tocopherol-.beta.-D-mannoside

Table 4: GC-MS Analysis of Phytochemical Compounds: Class and Biological Activity

Sl. no	Retention time	Name of the compound	Phytochemical class	Structure of compound	Biological activity
1	7.645	Isoamyl ethanoate	Ester	 C ₇ H ₁₄ O ₂	Antifungal, antibacterial, antiviral, anti-inflammatory & anti-haemolytic.
2	16.140	Anthranilic acid	Aromatic acid	 C ₇ H ₇ NO ₂	Antimicrobial, anti-inflammatory, analgesic, cytotoxic, and antioxidant activities
3	23.646	1H-Indole-2,3-dione	Indole alkaloid	 C ₈ H ₅ NO ₂	Antimicrobial, antidepressant, anticancer, anticonvulsant, and neuroprotective effects,
4	24.787	Trans-1,2-Diphenylcyclobutane	Cycloalkene	 C ₁₆ H ₁₆	Antimicrobial & antioxidant
5	25.580	2,3-Bis(1-methylallyl)pyrrolidine	Pyrrolidine alkaloid	 C ₁₂ H ₂₁ N	Antimicrobial activity & insecticidal
6	26.697	Neophytadiene	Diterpene	 C ₂₀ H ₃₈	Antimicrobial, antioxidant, anticancer, neuroprotective & cardiovascular.
7	27.120	Anthracene, 9-ethenyl-	Polycyclic aromatic hydrocarbon	 C ₁₆ H ₁₂	Antimicrobial, antioxidant, anti-inflammatory, cytotoxicity & anticancer,
8	28.189	3-Hydroxy-3-methyl-oxindole, Ac derivative	Oxindole alkaloid	 C ₁₁ H ₁₁ NO ₃	Antimicrobial, antioxidant, anti-inflammatory & anticancer
9	29.145	n-Hexadecanoic acid	Fatty acid	 C ₁₆ H ₃₂ O ₂	Antimicrobial, antioxidant & anticancer
10	32.009	Phytol	Diterpene alcohol	 C ₂₀ H ₄₀ O	Antimicrobial, antioxidant, anti-inflammatory, cytotoxicity, anticancer, neuroprotective & cardiovascular,
11	32.459	9,12,15-Octadecatrienoic acid (Z, Z, Z)-	Fatty acid (omega-3)	 C ₁₈ H ₃₀ O ₂	Antioxidant, anti-inflammatory, cytotoxicity, anticancer, neuroprotective & cardiovascular,
12	36.120	Cyclononasiloxane, octadecamethyl-		 C ₁₈ H ₅₄ O ₉ Si ₉	Industrial intermediate and cosmetic ingredient
13	37.652	Cyclohexane, 1,3,5-triphenyl-	Cycloalkane derivative	 C ₂₄ H ₂₂	Antioxidant & potential anticancer activity
14	39.385	(2,3-Diphenylcyclopropyl) methyl phenyl sulfoxide, trans-	Sulfoxide	 C ₂₂ H ₂₀ OS	Antimicrobial, antioxidant & anticancer
15	40.345	Tetracosamethyl - cyclododecasiloxane		 C ₂₄ H ₇₂ O ₁₂ Si ₁₂	No specific bioactivity
16	45.167	Trans-Indigo	Indole alkaloid	 C ₁₆ H ₁₀ N ₂ O ₂	Colorants, treatment of ulcerative colitis, antimicrobial, antioxidant & potential anticancer activity
17	48.426	Alpha, -Tocopherol- beta. -D-mannoside	Tocopherol	 C ₃₅ H ₅₈ O ₁₁	Antimicrobial, anti-inflammatory, anticancer & neuroprotective, antiviral & immunomodulatory

2,3-Diphenylcyclopropyl methyl phenyl sulfoxide isomers, with their unique stereochemistry, are of interest for their potential antioxidant, antimicrobial, and anticancer properties, as sulfoxide-containing compounds are explored for targeting cancer pathways. In addition, 9,12,15-Octadecatrienoic acid, an omega-3 fatty acid, is linked to cancer risk reduction and may support treatment efficacy. These compounds collectively suggest the sample may possess antioxidant, antimicrobial, and potential anticancer activities, making them relevant for further investigation in breast cancer and broader cancer research scenarios. The data will be given in Tables 3 and 4.

The analysis of the 17 compounds identified in the extract reveals several agents with strong biological activities related to various diseases and health benefits. Compounds such as 9,12,15-Octadecatrienoic acid ((Z, Z, Z)) exhibit antioxidant, anti-inflammatory (Nazir *et al.*, 2021), and proven anticancer properties, which can inhibit breast cancer cell proliferation, induce apoptosis, and potentially manage inflammatory conditions like arthritis (Yan *et al.*, 2024).

Cyclohexane, 1,3,5-triphenyl, exhibits antioxidant and possible anticancer activity, indicating that it may promote liver health, prevent oxidative stress in diabetes, and inhibit the growth of tumors in breast tissue. 2,3 Diphenyl cyclopropyl methyl phenyl sulfoxide exhibits antimicrobial (Kumar *et al.*, 2015), antioxidant, and anticancer properties that can fight infections, target breast cancer cells, and potentially control strains that are resistant to antibiotics (Havrankova *et al.*, 2020).

In addition to treating ulcerative colitis and acting as an antioxidant and anticancer agent, trans indigo may have immune-modulatory properties that are beneficial for the treatment of cancer, gut health, and autoimmune conditions like rheumatoid arthritis (Mizobuti *et al.*, 2022).

The antioxidant, anti-inflammatory, and anticancer properties of alpha tocopherol beta D protect cellular DNA, improve the effectiveness of chemotherapy against breast tumors, lessen the consequences of diabetes, and promote skin health (Reiter *et al.*, 2007).

CONCLUSION

Acetone proved to be an effective solvent for extracting diverse bioactive compounds from the *I. longiracemosa* plant, yielding 15% (w/w) crude extract rich in alkaloids, terpenoids, triterpenoids, fatty acids, and phenylpropanoids. The high recovery of phenylpropanoids and terpenoids supports the extract's strong antioxidant and potential anticancer properties. DPPH assay results showed concentration-dependent radical scavenging with an IC₅₀ of 9.25 µg/ml ($R^2 = 0.993$), excelling the standard antioxidant BHT (12 µg/ml; $R^2 = 0.9999$), indicating higher potency. GC-MS analysis revealed 25 peaks representing 17 compounds, including α-linolenic acid, n-hexadecanoic acid, phytol, neophytadiene, indole, oxindole, trans-indigo, and α-tocopherol β-D-mannoside, many known for antioxidant, anti-inflammatory, and anticancer effects. These findings suggest the extract's therapeutic relevance as a natural source of multifunctional bioactive molecules.

Overall, the extraction efficiency, strong DPPH radical scavenging with an IC₅₀ lower than BHT, and the phytochemical profile defined by GC-MS suggest that this acetone extract is a rich source of bioactive molecules with significant antioxidant potential and possible relevance to cancer, especially breast cancer. However, antioxidant assays like DPPH represent just an initial *in vitro* screening step and cannot predict *in vivo* effectiveness or safety on their own. *In vitro* cytotoxicity and mechanism assessments in breast cancer cells, examining apoptosis and cell cycle control, followed by *in vivo* testing in a suitable mammary cancer model, confirmed the anticancer potential indicated by the current results. Additionally, isolating the most active components through fractionation and bioactivity-guided methods, combined with molecular pathway analysis, will clarify the roles of these phytochemicals and may ultimately lead to new plant-based treatments or lead compounds for breast cancer prevention and therapy.

Acknowledgements

The authors declare that no funds, grants, or other support were received during the preparation of this manuscript. The GC-MS data presented in this study are unpublished and part of the current research.

Conflict of interest

The authors declare no competing interests.

Author contributions

Shabna. K: Writing – review & editing, Writing- original draft, Methodology, Investigation, Data curation, Conceptualization. Dr Arun Kumar G: Visualization, Validation, Supervision, and Investigation.

Data Availability Statement: Not applicable.

Correspondence and requests for materials should be addressed to **K. Shabna**

Orcid id

Author 1: SHABNA. K ([0009-0009-3477-522X](https://orcid.org/0009-0009-3477-522X))

Author 2: Arun Kumar G ([0000-0002-5984-9899](https://orcid.org/0000-0002-5984-9899))

Reprints and permissions information is available at

<https://www.ijlsci.in/reprints>

REFERENCES

- Altemimi A, Lakhssassi N, Baharlouei A, Watson DG, Lightfoot DA (2017) Phytochemicals: Extraction, isolation, and identification of bioactive compounds from plant extracts, *Plants*, 6(4) Article 42. <https://doi.org/10.3390/plants6040042>
- Blois MS (1958) Antioxidant determinations by the use of a stable free radical, *Nature*, 181(4617) 1199–1200.
- Bray F, Ferlay J, Soerjomataram I, Siegel RL, Torre LA, Jemal A (2018) Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries, *CA Cancer J Clin*, 68(6) 394–424. <https://doi.org/10.3322/caac.21492>
- Cragg GM, Newman DJ, Snader KM (2018) Natural products in drug discovery and development, *J Nat Prod*, 81(3) 513–518. <https://doi.org/10.1021/acs.jnatprod.9b01285>
- Dehelean CA, Marcovici I, Soica C, Mioc M, Coricovac D, Iurciuc S, Cretu OM, Pinzaru I (2021) Plant-derived anticancer compounds as new perspectives in drug discovery and alternative therapy, *Molecules*, 26(4) 1109. <https://doi.org/10.3390/molecules26041109>
- Havrankova E, Calkovska N, Padrtova T, Csollei J, Opatrilova R, Pazdera P (2020) Antioxidative activity of 1,3,5-triazine analogues incorporating aminobenzene sulfonamide, aminoalcohol/phenol, piperazine, chalcone, or stilbene motifs, *Molecules*, 25(8) 1787. <https://doi.org/10.3390/molecules25081787>
- Hethelyi E, Tetenyi P, Dabi E, Danos B (1987) The role of mass spectrometry in medicinal plant research, *Biomed Environ Mass Spectrom*, 14(11) 627–632.
- Hubschmann HJ (2015) Handbook of GC–MS: Fundamentals and applications (3rd ed.), Wiley-VCH. <https://doi.org/10.1002/9783527674305>
- Kumar S, Kumar V, Prakash O (2009) Antipyretic activity of *Indigofera longiracemosa*, *J Ethnopharmacol*, 126(3) 473–478.
- Kumar S, Kumar V, Kumar R, Singh D (2015) Extraction, isolation, and characterization of bioactive compounds from medicinal plants: A review, *J Pharmacogn Phytochem*, 4(3) 28–35.
- Mizobuti DS, da Rocha GL, da Silva HNM, Covatti C, de Lourenco CC, Pereira ECL, Salvador MJ, Minatel E (2022) Antioxidant effects of bis indole alkaloid indigo and related signalling pathways in the experimental model of Duchenne muscular dystrophy, *Cell Stress Chaperones*, 27(4) 417–429. <https://doi.org/10.1007/s12192-022-01282-0> (doi.org in Bing)
- Mukherjee PK, Wahile A (2018) Integrated approaches towards drug development from Ayurveda and other Indian systems of medicine, *J Ethnopharmacol*, 211 259–269. <https://doi.org/10.1016/j.jep.2017.09.024>
- Nazir N, Zahoor M, Uddin F, Nisar M (2021) Chemical composition, in vitro antioxidant, anticholinesterase, antimicrobial, anti-urease and antileishmanial activities of *Calligonum polygonoides* L, *Sci Rep*, 11(1) 4369. <https://doi.org/10.1186/s12906-021-03228-y>
- Pan SY, Zhou SF, Gao SH, Yu ZL, Zhang SF, Tang MK, Sun JN, Ma DL, Han YF, Fong WF, Ko KM (2013) New perspectives on how to discover drugs from herbal medicines: CAM's outstanding contribution to modern therapeutics, *Evid Based Complement Alternat Med*, 2013 627375. <https://doi.org/10.1155/2013/627375>
- Patel DK, Kumar R, Laloo D, Hemalatha S (2013) Antioxidant activity of *Indigofera longiracemosa* roots, *J Ethnopharmacol*, 146(3) 855–861.
- Reiter E, Jiang Q, Christen S (2007) Anti-inflammatory properties of α - and γ -tocopherol, *Mol Aspects Med*, 28(5–6) 668–691. <https://doi.org/10.1016/j.mam.2007.01.003>
- Sebaugh JL (2011) Guidelines for accurate EC50/IC50 estimation, *Pharm Stat*, 10(2) 128–134. <https://doi.org/10.1002/pst.426>
- Siegel RL, Miller KD, Jemal A (2020) Cancer statistics, 2020, *CA Cancer J Clin*, 70(1) 7–30. <https://doi.org/10.3322/caac.21590>
- Singh GN, Singh P, Kumar P (2012) Wound healing activity of *Indigofera longiracemosa* leaves, *J Ethnopharmacol*, 143(2) 551–558.
- Sung H, Ferlay J, Siegel RL, Laversanne M, Soerjomataram I, Jemal A, Bray F (2021) Global cancer statistics 2020: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries, *CA Cancer J Clin*, 71(3) 209–249. <https://doi.org/10.3322/caac.21660>
- Thangadurai D (1998) Ethnomedicinal plants used as an antidote for poisonous bites among the tribal of the southern Western Ghats, India, *Proc Natl Conf Med Plant Res*, Calcutta, West Bengal.
- Uma G, Balasubramaniam V (2012) GC–MS analysis of *Nothapodytes nimmoniana* Mabblerly leaves, *J Chem Pharm Res*, 4(9) 4417–4419.
- Wagner H, Bladt S, Zgainski EM (1984) Plant drug analysis: A thin layer chromatography atlas, Springer-Verlag, Berlin. <https://doi.org/10.1007/978-3-642-71474-9>

- Weissenstein U, Kunz M, Urech K, Baumgartner S (2014) Interaction of standardized mistletoe (*Viscum album*) extracts with chemotherapeutic drugs regarding cytostatic and cytotoxic effects in vitro, *Evid Based Complement Alternat Med*, 2014 1–9. <https://doi.org/10.1155/2014/>
- Yan H, Zhang S, Yang L, Jiang M, Xin Y, Liao X, Li Y, Lu J (2024) The antitumor effects of α -linolenic acid, *Nutrients*, 14(3) 260. <https://doi.org/10.3390/jpm14030260>.

© The Author(s) 2026

Publisher's Note

IJLSCI remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.