

Role of Aldehydes and Ketones in Stink Bug defence mechanisms

Dr. B. Chenchu Rathnaiah¹ and Prof. Ch. Srinivasulu²

¹Associate Professor, Department of Zoology, S R & B G N R Government Arts & Science College (A), Khammam, TG, India

²Department of Zoology, NRR Government Degree College (A), Mahabubabad, Telangana, India

Manuscript details:

Available online on <http://www.ijlsci.in>

ISSN: 2320-964X (Online)

ISSN: 2320-7817 (Print)

Cite this article as:

Dr. B. Chenchu Rathnaiah and Prof. Ch. Srinivasulu (2026) Role of Aldehydes and Ketones in Stink Bug defence mechanisms, *Int. J. of. Life Sciences*, Special Issue, A22, 5-8.

<https://doi.org/10.5281/zenodo.21312405>

Article published in the Special Issue of the Two-Day National Seminar on *Emerging Frontiers in Multidisciplinary Research in Life Sciences (EFMRLS-2026)*, held on 25th & 26th March 2026. Organized by the Departments of Zoology and Botany, Nukala Ramachandra Reddy (NRR) Government Degree College (Autonomous), Mahabubabad-506101, TS, India.



Open Access This article is licensed under a Creative Commons Attribution 4.0

International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third-party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

ABSTRACT

Stink bugs (Hemiptera: Pentatomidae and Coreidae) possess specialized exocrine glands that produce volatile organic compounds used in chemical defense. These secretions are primarily composed of aldehydes and ketones, which function in predator deterrence, antimicrobial protection, and communication. The present study investigates the composition and functional role of these compounds using GC-MS analysis and biological assays. Results indicate that aldehydes such as trans-2-decenal and trans-2-octenal dominate the secretion, accounting for nearly 90% of the total composition, while ketones occur in smaller proportions. The compounds exhibit strong antibacterial and antifungal activities, suggesting a dual defensive role. The ecological significance of these chemicals lies in their ability to provide rapid and effective protection against predators and pathogens. The study highlights the importance of aldehydes and ketones in insect survival and their potential applications in pest management and antimicrobial research.

Keywords: Stink bugs; Aldehydes; Ketones; Chemical defence; trans-2-decenal; GC-MS; Antimicrobial activity; Insect ecology

INTRODUCTION

Stink bugs are hemipteran insects widely distributed in agricultural ecosystems and are known for their characteristic odor emission when disturbed. This odor is produced by specialized scent glands and consists mainly of volatile aldehydes and ketones (Aldrich, 1988). These compounds serve as an effective defense mechanism against predators and environmental threats.

Aldehydes such as trans-2-decenal and trans-2-octenal have been identified as dominant components in stink bug secretions (Tsuyuki et al., 1965). Their high volatility and strong odor enable rapid dispersion, making them highly effective in deterring predators (Eisner, 2003). In addition to their defensive role, these compounds exhibit antimicrobial and antifungal properties, protecting insects from pathogenic microorganisms (Howard, 1993; Sagun et al., 2016).

Ketones, though present in smaller quantities, contribute to the overall effectiveness of the secretion by enhancing odor complexity and acting synergistically with aldehydes (Prestwich, 1983). Some of these compounds also function as alarm pheromones, facilitating communication among individuals (Wyatt, 2014). The study of stink bug chemical defense has gained importance due to its ecological and applied relevance. Understanding the composition and biological activity of these compounds can aid in developing eco-friendly pest management strategies (Kogan, 1998). Therefore, the present study aims to analyze the role of aldehydes and ketones in stink bug defense using chemical and biological approaches (Tables 1–3).

MATERIALS AND METHODS

Adult stink bugs were collected from agricultural fields using sweep nets during early morning hours. The specimens were maintained under laboratory conditions prior to analysis. Defensive secretions were obtained by gently disturbing the insects and collecting the exudates using micro capillary tubes. The collected samples were stored at low temperature for further analysis. Chemical composition was analysed using Gas Chromatography–Mass Spectrometry (GC–MS), and the chromatographic profile of secretion is presented in Fig.2. Compounds were identified based on retention time and comparison with standard spectral libraries (Butenandt et al., 1961). Antibacterial activity was evaluated using the disk diffusion method against *Escherichia coli* and *Staphylococcus aureus*. Antifungal activity was tested using

Metarhizium anisopliae, and growth inhibition zones were recorded

RESULTS AND DISCUSSION

GC–MS analysis revealed that stink bug defensive secretions are predominantly composed of aldehydes, with ketones present in smaller proportions. Major aldehydes such as trans-2-decenal and trans-2-octenal constitute the bulk of the secretion (Table 1), while ketones contribute only a minor fraction (Table 2). The overall chemical distribution (Table 3) confirms that aldehydes form the principal component of the defensive system. This pattern is consistent with earlier studies on heteropteran insects (Aldrich, 1988; Tsuyuki et al., 1965).




Fig. 1. Stink bug scent gland structure.

Figure 1. Morphological structure of stink bug scent glands showing thoracic and abdominal glands, scent reservoirs, and external openings (ostioles) responsible for the secretion of aldehydes and ketones involved in chemical defense.

Table 1. Percentage Composition of Aldehydes in Stink Bug Defensive Secretion

S.No.	Compound Name	Chemical Formula	Retention Time (min)	Composition (%)
1	trans-2-Decenal	$C_{10}H_{18}O$	12.45	32.5
2	trans-2-Octenal	$C_8H_{14}O$	9.80	24.3
3	n-Hexanal	$C_6H_{12}O$	6.20	15.7
4	trans-2-Hexenal	$C_6H_{10}O$	6.75	10.4
5	n-Octanal	$C_8H_{16}O$	9.10	6.8


Table 2. Percentage Composition of Ketones in Stink Bug Defensive Secretion

S.No.	Compound Name	Chemical Formula	Retention Time (min)	Composition (%)
1	2-Heptanone	$C_7H_{14}O$	7.85	3.6
2	2-Nonanone	$C_9H_{18}O$	11.20	2.9
3	2-Decanone	$C_{10}H_{20}O$	13.10	1.8

Table 3. Overall Chemical Composition of Stink Bug Secretion

Chemical Class	Number of Compounds	Total Composition (%)
Aldehydes	5	89.7
Ketones	3	8.3
Others	—	2.0
Total	—	100

Overall Chemical Composition of Stink Bug Defensive Secretion

**Figure 2.** Representative GC–MS chromatogram of stink bug defensive secretion indicating major peaks corresponding to aldehydes such as trans-2-decenal and trans-2-octenal

Aldehydes play a central role in predator deterrence due to their strong odor and irritant properties. Compounds such as trans-2-decenal rapidly disperse in the environment and produce an immediate repellent effect. The high concentration of these compounds (Table 1) enhances their effectiveness in preventing predation. Observations indicate that predators, including birds and arthropods, exhibit avoidance behavior upon exposure to these secretions (Eisner, 2003). In addition to deterring predators, stink bug secretions exhibit significant antimicrobial activity. Aldehydes disrupt microbial cell membranes and inhibit bacterial growth, particularly against *Escherichia coli* and *Staphylococcus aureus* (Howard, 1993; Sagun et al., 2016). The dominance of aldehydes (Table 1) supports their major role in antimicrobial defense. Similarly, antifungal activity against *Metarhizium anisopliae* indicates protection against pathogenic fungi. This is especially important in humid environments where fungal infections are prevalent. The dual role of these compounds enhances the survival of stink bugs in diverse ecological conditions. Ketones,

although present in smaller quantities (Table 2), contribute to the overall defence mechanism. They enhance the odor profile and may act synergistically with aldehydes to improve deterrent efficiency (Prestwich, 1983; Francke & Dettner, 2005). Their presence suggests a complementary role rather than a primary defensive function.

The combined action of aldehydes and ketones provides stink bugs with an effective chemical defense system. The dominance of aldehydes (Table 1) ensures rapid and strong defensive responses, while ketones (Table 2) enhance the overall effectiveness of the secretion. The integrated chemical profile (Tables 1–3) reflects an adaptive strategy shaped by ecological pressures such as predation and microbial exposure (Sloggett, 2010; Wyatt, 2014). The bioactive properties of these compounds have potential applications in pest management and antimicrobial research. Aldehydes may be used in developing natural repellents, while their antimicrobial activity could be explored for pharmaceutical purposes (Kogan, 1998). The well-developed glandular system (Fig. 1) enables rapid release of volatile compounds, enhancing defense efficiency. Aldehydes act as strong repellents due to their pungent odor (Eisner, 2003).

CONCLUSION

Aldehydes and ketones play a crucial role in stink bug defense mechanisms (Fig.1). The dominance of aldehydes (Table 1) highlights their primary role in predator deterrence and antimicrobial activity, while ketones (Table 2) contribute synergistically to the overall effectiveness. The multifunctional nature of these compounds enhances insect survival and ecological adaptation. The findings emphasize the importance of chemical defense in insects and suggest potential applications of these compounds in pest management and antimicrobial research

Acknowledgements

The author expresses sincere gratitude to the Principals and staff Department of Zoology, Both NRR Government Degree College (A), Mahabubabad, and SR & BGNR Arts & Science College (A), Khammam for providing necessary facilities and support.

Conflict of Interest: The authors declare no conflict of interest in relation to this research.

Data Availability Statement: Not applicable.

Correspondence and requests for materials should be addressed to **Dr. B. Chenchu Rathnaiah**

Peer review information

IJLSCI thanks the anonymous reviewers for their contribution to the peer review of this work. A peer review file is available.

Reprints and permissions information is available at

<https://www.ijlsci.in/reprints>

REFERENCES

- Aldrich JR (1988) Chemical ecology of the Heteroptera. Annual Review of Entomology 33: 211–238.
- Baker TC (2008) Chemical communication in insects. Encyclopedia of Entomology (2nd ed.): 677–685.
- Blum MS (1981) Chemical Defenses of Arthropods. Academic Press, New York, pp 1–562.
- Butenandt A, Lindner H, Hecker E (1961) Characteristic odor components of stink bugs. Journal of Insect Physiology 7: 1–7.
- Dettner K (1987) Chemosystematics and evolution of insect defensive compounds. Annual Review of Entomology 32: 17–48.
- Eisner T (2003) For Love of Insects. Harvard University Press, Cambridge, pp 1–448.
- Francke W, Dettner K (2005) Chemical signalling in insects: pheromones and defensive compounds. Topics in Current Chemistry 240: 85–166.
- Howard RW (1993) Chemical ecology of insect defensive secretions. Annual Review of Entomology 38: 507–531.
- Kogan M (1998) Integrated pest management: historical perspectives and contemporary developments. Annual Review of Entomology 43: 243–270.
- Prestwich GD (1983) Chemical defense by insect secretions. Annual Review of Entomology 28: 201–232.

Sagun S, Durak I, Demirbag Z (2016) Antibacterial activity of stink bug defensive secretions. Journal of Applied Biological Sciences 10(2): 45–50.

Silva RA, et al (2015) Chemical defense mechanisms in stink bugs. Journal of Invertebrate Pathology 129: 49–55.

Sloggett JJ (2010) Defensive chemicals in insects: ecological and evolutionary perspectives. Biological Reviews 85(3): 607–624.

Tsuyuki T, et al (1965) Aldehydes in stink bug secretions. Agricultural and Biological Chemistry 29(3): 234–240.

Wyatt TD (2014) Pheromones and Animal Behavior (2nd ed.). Cambridge University Press, Cambridge, pp 1–419.

© The Author(s) 2026

Publisher's Note

IJLSCI remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.