



# Metabolic impact on lipid content of *Oreochromis mossambicus* after lead nitrate intoxication

Chavan PN<sup>1</sup> and Mali RP<sup>2</sup>

<sup>1</sup>Department of Zoology, S.G.B. Mahavidyalaya, Purna, MS, India

<sup>2</sup>Indira Gandhi Mahavidyalaya, Nanded, MS, India

\*Corresponding author email- [pnychavan.2009@rediffmail.com](mailto:pnychavan.2009@rediffmail.com)

## Manuscript details:

Received: 01.11.2020  
Accepted: 10.11.2020  
Published: 30.12.2020

## Cite this article as:

Chavan PN and Mali RP (2020) Metabolic impact on lipid content of *Oreochromis mossambicus* after lead nitrate intoxication, *Int. J. of Life Sciences*, 2020; 8 (4):774-776.

Available online on <http://www.ijlsci.in>  
ISSN: 2320-964X (Online)  
ISSN: 2320-7817 (Print)



Open Access This article is licensed under a Creative Commons

Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

## ABSTRACT

The heavy metal pollutants released into natural aquatic medium by anthropogenic activities which later interact with aquatic organisms. The contamination of water through heavy metal is potential problem for aquatic organism. The present investigation was the event on estimation of lipid content from different tissues of *Oreochromis mossambicus* after exposed to lead nitrate intoxication. In present study the impact of lead nitrate on *Oreochromis mossambicus* has been studied for 15 days at 96 hours exposure and compared with control set. The observation and results discussed in detail.

**Keywords:** Lipid, *Oreochromis mossambicus*, Lead Nitrate

## INTRODUCTION

Among the different habitats aquatic environment is the major target of pollution. Most of the heavy metals are natural constituents of the aquatic environment. The pollution of water is the greatest and most problematic source due to industrialization. The impact of pollution on aquatic biota and ecosystem, human health is a recent international issue which creates the environmental disturbances. Most of the heavy metals are natural constituents of the aquatic environment, some of them are biologically essential and some metals like lead, cadmium and mercury are hazardous to aquatic biota (Mali, 2002). The main source of water pollution is domestic sewage, pesticides, fertilizers and industrial effluents which pollutes water resources (Maruthanayagam and Sharmila, 2004). The contamination of water through heavy metal is potential problem for aquatic biota. The higher concentration of heavy metal leads to alteration in physico-chemical and biological properties of water and can cause hazardous effect on aquatic biota (Jagadeesan *et al.*, 2001).

The nutritive and medicinal values of fish have been recognized from immortal time. The heavy metal concentration in different tissues of fish enters into human beings through food chain (EL-Shehawi *et al.*, 2007).

The impact of heavy metal directly affects the biochemical constituents of aquatic organisms. Hence it is necessary to understand the significance of these variations in the organic compound of the tissues. The present study was carried out to investigate the lipid content in different tissues of fish after exposure to lead nitrate.

## MATERIAL AND METHOD

The fish *Oreochromis mossambicus* were collected from Godavari River Nanded (M.S) with the help of local fisherman. The fishes were kept in glass aquarium and fed with slice of tubifex. They were acclimatized in

laboratory condition for ten days. Then these fishes are divided in two groups. The group A for control set and group B for sub lethal concentration of lead nitrate of 10 ppm. The tissue like liver, muscle and gill were selected for experimentation. The total lipid content was done by Methanol-Chloroform method by Folch *et al.*, (1957).

## RESULT AND DISCUSSION

Lead nitrate is one of the most common heavy metal that affect adversely on freshwater fishes. The results of different tissues of fish are presented in Figure 1.




Figure 1: Lipid content in liver, muscle and gill of *Oreochromis mossambicus* after lead nitrate intoxication.

There was a remarkable change in the total lipid content in fish. In control set the amount of lipid in liver 14.50, while the treated fish showed 9.31 lipid mg/gm wet wt. of tissue. The muscle of control set showed 17.22, whereas treated fish showed 14.44 lipid mg/gm wet wt. of tissue. The gill of control set showed 10.02 and treated fish 8.40 lipid mg/gm wet wt. of tissue. The present investigation showed decline in the lipid content of fish as compared to control set. Similar results were observed in fresh water fingerling *Labeo rohita* (Raja and Puvaneshwari, 2017). Leela *et al.*, (2000) showed significance decrease in total lipid content of muscle, liver and gill of *Tilapia mossambica*

under the stress of phosalone. The decrease in lipid content in the different tissues suggested that the lipid have been channalised to meet the metabolic demand for extra energy need to overcome the toxic stress. Katti and Sathyanesan (1983) showed that decreased cholesterol and lipid level in brain, testis and ovary of *Clarius batrachus* exposed to 5ppm of lead nitrate for 150 days. In the present investigation there was decrease in the lipid content of all the tissues of fresh water fish after lead nitrate intoxication, similar results were obtained in *cirrhinus mrigala* after acute and chronic exposure of heavy metals (Bhilave *et al.*, 2008).

**Conflicts of Interest:** The author declares no conflict of interest

## REFERENCES

- Bhilave MP, Muley DV and Deshpande VY (2008) Biochemical changes in the fish *Cirrhinus mrigala* after acute and chronic exposure of heavy metals.
- El-Shehawi AM, Ali FK and Seehy MA (2007) Estimation of water pollution by genetic biomarkers in tilapia and cat fish species shows species site interaction, *Afr. J. Biotech.*, 6840-846.
- Folch J, Lees M and Sloane-Stanley GH (1957) A simple method for the isolation and purification of total lipids from animal tissue. *J. Biol. Chem.* 226: 497- 509.
- Jegadeesan G, Jebanesan A and Mathivanan A (2001) In vivo recovery of organic Constituents in gill tissue of *Labeo rohita* after exposure to sub lethal concentrations of mercury. *J. Exp. Indellerria* 3: 22-29.
- Katti SR and Sathyanesan AG (1983) Lead nitrate induced changes in lipid and cholesterol level in the fresh water fish *Clarius battachus*. *Toxicol. Lett.*, 19 (1-2); 93-96.
- Leela Shiva Parvathi M, Chander Sekhara Reddy D, Nadamuni Chetty A (2000) In vivo recovery and long term effect of Phosalone on total lipid and triglyceries in fresh water fish *Tilapia mossambica* (1 Peters) *Poll. Res.* 19 (3): 345-351.
- Mali RP (2002) Studies on some aspects of physiology of freshwater female crab *Barytelphusa guerini* with special reference to inorganic pollutants. Thesis submitted to S.R.T.M.U., Nanded.
- Muruthanayagam C and Sharmila G (2004) Hemato-Biochemical Variations induced by the pesticide monocrotophos in *Cyprinus carpio* during the exposure and recovery periods *Nat. Environ, Poll. Tech.* 3: 49991-494.
- Raja P and Puvaneshwari S (2017) The heavy metal lead nitrate toxicity effect on biochemical alteration in fresh water fingerlings *Labeo rohita*, (Hamilton, 1882) *International Journal of Advanced science and research* 2: 58-63.