



# Effects of heavy metals on the ventilation and opercular movement in *C. gachua* and *C. reba*

Mandip Kumar Roy<sup>1\*</sup> and Akriti Rani<sup>2</sup>

<sup>1</sup>Department of Zoology, T.P. Varma College, Narkatiaganj, West Champaran (Bihar) – 845455

<sup>2</sup>Research Associate, J.P University, Chhapra (Bihar)

\*Corresponding author email: Email: [drmandiproym@gmail.com](mailto:drmandiproym@gmail.com)

## Manuscript details:

Received: 01.07.2022  
Revised: 22.07.2022  
Accepted: 26.08.2022  
Published: 30.09.2022

## Cite this article as:

Mandip Kumar Roy and Akriti Rani (2022) Effects of heavy metals on the ventilation and opercular movement in *C. gachua* and *C. reba*, *Int. J. of Life Sciences*, 10 (3): 233-238.

Available online on <http://www.ijlsci.in>

ISSN: 2320-964X (Online)

ISSN: 2320-7817 (Print)



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other thirdparty material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

## ABSTRACT

Several heavy metals which are present as pollutants in the water bodies have been established a toxic substances to fishes. The present study has been aimed to regulate their entry in the various water bodies so that it can be ensured that they may not be adversely affected in any way. It has been found that the effects of heavy metals in fish can be variable to a greater extent. Apart from causing death either directly or due to starvation by destruction of food organism, many heavy metals have shown to affect the survival behaviour, growth, reproduction, physiology biochemistry and pathology of fish with evidence of tissue damage. The present study deals with the opercular frequencies and respiratory movement. It was found that their frequencies and distress were increased 2-4 times more than the control fishes. The opercular frequency/minute were observed in two different habit fishes at different concentration to two heavy metals i.e.  $\text{CuSO}_4$  and  $\text{K}_2\text{Cr}_2\text{O}_7$  pollutants in relation to exposure periods. At the initial stage of exposure, both fishes i.e. *C. gachua* and *C. reba* exhibited similar result and no significant affect were noticed. However in case of *C. gachua*, the opercular frequency was found in increasing there from 40mg/l to 265.4 mg/l level but suddenly and abruptly declined at 517.9mg/l level up to 72 hour of exposure in both pollutants Another test fish *C. reba* exhibited some different affects i.e increasing trend from 24 hour exposure and started decreasing at 9b hr. of exposure which was not abrupt like *C. gachua* rather in or systematic manner except in case of  $\text{CuSO}_4$  pollutants of 88mg/l level of concentration where it exhibited  $46 \pm 2$ . The initial increase in the opercular movement followed by a gradual decrease which becomes rapid just prior to death. These development depended mainly on the concentration limit and exposure period and normally this is concentration dependent Further the movement of *C. gachua* and *C. reba* towards water volume increased in almost all concentration to meet their  $\text{O}_2$  requirement but it did not materialize in lethal concentration.

**Keywords:** Heavy metals, *C.gachna*, *C. reba*, ventilation, opercular movement,  $\text{CuSO}_4$ ,  $\text{K}_2\text{Cr}_2\text{O}_7$

## INTRODUCTION

Aquatic pollution has become a global problem in recent years. Extensive industrialization has influenced the quality of water all over the world. Heavy metals are recognized as one of the most hazardous pollutants and toxic to many organisms. The natural aquatic systems extensively contaminate with heavy metals released from domestic, industrial and other man-made activities (Vutukuru, 2003 Monazaki et. al. 2015).

Fishes are widely used to evaluate the health of aquatic ecosystem. They respond sensitivity to an increased concentration of contaminants in water. Heavy metals at high concentrations can cause hazardous effects to metabolic, physiological and biochemical system of fishes (Health. 1987).

Among the heavy metals, cadmium, lead, mercury, copper, zinc, chromium and nickel are comparatively notorious toxicants and most of their compounds are water soluble and non-degradable zinc and copper which are used in agriculture works and are known as essential element in animals and plants metabolism are also reported toxic to fishes (Singh and Singh 1979).

The initial increase in the opercular movement followed by a gradual decrease which becomes rapid just prior to death. These development depended mainly on the concentration limit and exposure period and normally this is concentration dependent further the movement of *C. reba* and *C. gachua* towards water volume increased in almost all concentration to meet their oxygen requirement but it did not materialize in lethal concentration.

The present study deals with the opercular frequencies and respiratory distress in the presence of undesirable concentration of  $\text{CuSO}_4$  and  $\text{K}_2\text{Cr}_2\text{O}_7$ . It is an attempt towards the aim of reducing the level of pollution to those which can at least be tolerated and beneficial to the fishes and other aquatic biota and also for maintaining healthy stock of fish.

## MATERIALS AND METHODS:

Mature specimen of healthy and alive *Channa gachua* commonly known as "Changa" of  $36.5 \pm 2.5$  gm, weight

groups and *Cirrhinus reba* commonly known as "Reba" of both sexes collected from the aquatic resources situated in the vicinity of Motihari town and brought to the laboratory, where they washed for few minutes in 0.1% aqueous potassium permanganate solution followed by several changes of fresh water to remove any dermal infection present.

During acclimation and experimental periods, fishes were provided with "Fish tone" (an artificial food), tubifex and chopped goat's liver and alternate day respectively two to three hours prior to the change of water and chronic exposure except bio-assay test, during which fishes were not provided food twenty four hours prior to start the experiment to the end of the experiment (i.e. up to 96 hours).

*Bioassay test:* Static bioassay tests for determining the acute toxicity or median lethal concentrations ( $\text{LC}_{50}$ ) value at 24, 48, 72 and 96 hr of exposures to the selected metallic salts ( $\text{CuSO}_4$  and  $\text{K}_2\text{Cr}_2\text{O}_7$ ) individually were patterned following the method described by Doudoroff et al (1951) and APHA, (1995).

The opercular frequencies or ventilation rates of five fish of each concentration and the controls at selected periods of exposures (i.e. 24, 48, 72 and 96 hr of exposure) were counted by visual observation/using magnifying glass for 5-10 minutes and average value per minutes/fish were calculated and recorded.

## RESULTS AND DISCUSSION:

During the experimental period no death was recorded in the test level and at lowest concentration of heavy metals. Both fish exhibited characteristics but identical behaviorable changes depending upon the concentrations and exposure period. Their increased opercular movement initially reflected the restlessness evidence of difficulty in respiration of the fishes was apparent because they surfacing. The poison symptomology usually started before three to four hrs. of death as observed by loss of equilibrium, convulsion and erratic jerky body action. The fishes gradually became lethargic followed by paralysis and finally they settled down at the bottom of the container. The respiratory distress along with opercular frequency was seen to be increased by 2-4 times more than the control fish.

**Table 1:** Opercular frequency per minute in relation to exposure period and concentrations of the heavy metal during acute exposure in *Channa gachua*.

Heavy Metal	Concentration (ppm)	Starting point	Hour of exposure			
			24	48	72	96
Test	00	38±4	36±3	38±3	35±3	38±3
CuSO <sub>4</sub>	40	38±4	43±3	41±3	36±3	27±3
	60	38±4	41±3	42±2	32±2	27±3
	88	38±4	38±3	41±2	35±2	29±2
	127.4	38±4	49±2	41±1	33±2	29±2
	182.2	38±4	51±1	45±4	36±1	28±1
	285.1	38±4	50±4	37±3	28±4	20±2
	265.1	38±4	50±3	29±2	19±3	-
	517.1	38±4	35±2	24±2	-	-
	0	38±4	35±2	39±2	38±2	39±2
K <sub>2</sub> Cr <sub>2</sub> O <sub>7</sub>	40	38±4	36±2	41±2	38±2	34±2
	60	38±4	51±4	45±4	35±4	31±4
	88	38±4	41±5	42±5	41±5	28±5
	127.4	38±4	40±2	39±2	74±2	24±2
	182.2	38±4	52±4	45±2	33±4	18±4
	258.1	38±4	51±2	32±2	19±2	11±2
	265.4	38±4	28±3	20±3	18±3	-
	517.9	38±4	21±2	18±2	-	-

± is standard error of 4 observation

**Table 2:** Opercular frequency per minute in relation to exposure period and concentrations of the heavy metal during acute exposure in *C. reba*.

Heavy Metal	Concentration (ppm)	Starting point	Hour of exposure			
			24	48	72	96
Test	00	38±2	38±3	39±3	37±3	37±3
CuSO <sub>4</sub>	40	38±4	44±3	45±3	38±3	39±3
	60	38±4	42±2	44±2	46±2	36±2
	88	38±4	41±2	39±2	38±2	31±2
	127.4	38±4	48±2	42±2	35±2	29±2
	182.2	38±4	52±3	46±3	34±3	28±3
	285.1	38±4	42±2	41±2	31±2	27±2
	265.1	38±4	45±2	40±2	29±2	26±2
	517.1	38±4	47±3	42±3	29±3	25±2
	0	38±4	41±2	37±1	38±2	39±2
K <sub>2</sub> Cr <sub>2</sub> O <sub>7</sub>	40	38±4	39±3	39±3	37±3	37±3
	60	38±4	38±3	37±3	39±3	37±3
	88	38±4	39±3	36±3	38±3	35±3
	127.4	38±4	39±2	38±2	36±2	34±2
	182.2	38±4	40±3	38±3	39±3	34±3
	258.1	38±4	41±2	40±2	37±2	33±2
	265.4	38±4	43±3	39±3	36±3	30±3
	517.9	38±4	28±3	37±3	35±3	29±3

The opercular frequency per minute were observed in both fishes at different concentration levels to both metal pollutants in relation to exposure periods. The frequency were studied at 24, 48, 72 and 96 hrs.

In case of *C. gachua* when exposed to 24, 48, 72 and 96 hr. at different concentration level to CuSO<sub>4</sub> the opercular frequency was found 38 ±2 at starting point and was found to be increased from 40mg/l

concentration to 265.4 mg/l level but suddenly and abruptly declined at 517.9mg/l level. The opercular movement of *C. gachua* when exposed to  $K_2Cr_2O_7$  at different level at selected hr. of exposures revealed more abruptly declining trend at 265.4 and 517.9mg/l concentration level. At 48 hr. of exposure almost similar trend was noticed when exposed to both pollutants but decreasing level was starting at  $38 \pm 3$  and  $39 \pm 2$  after increasing level to some concentration level, abruptly decreased to  $29 \pm 2$  and  $24 \pm 2$  and  $20 \pm 3$  and  $18 \pm 2$  respectively to both  $CuSO_4$  and  $K_2Cr_2O_7$ . At 72 hr. of exposure to both pollutants, the opercular frequency was observed to be in the increasing, decreasing and increasing trends abruptly decreased to  $19 \pm 3$  and  $18 \pm 3$  respectively at 265.4mg/l. level concentration and no movement was seen at 517.9 mg/l concentration to both pollutants (Table-1).

The opercular movement of *C. gachua* to both pollutants at 96 hrs of exposure at different concentration level exhibited decreasing trend except in case of  $CuSO_4$  exposure at 88 and 258.1 mg/l level concentration. This frequency was finally come to half at 265.4mg/l concentration level.

The opercular frequency per minute in relation to exposure periods and concentration of both heavy metals with percent change in case of *C. gachua* has been shown in Table-1.

In case of *C. reba* the opercular frequency were too observed. It exhibited increasing, decreasing and increasing trend and showing nothing extra action.

At 24 hrs. of exposure to  $CuSO_4$  pollutants exhibited increasing decreasing and increasing trend and finally the opercular movement was found to be  $47 \pm 3$  at 517.9 mg/l level of concentration. Similar trend was noticed to  $K_2Cr_2O_7$  pollutants except that it abruptly decreased  $28 \pm 3$  at 517.9 level.

At 48 hr. of exposure to  $CuSO_4$  the opercular frequency was observed to be in the similar fashion as per the result at 24 hr. of exposure above but at  $K_2Cr_2O_7$  pollutants the increasing and decreasing level of opercular frequency does not show any significant variation rather it moves in between  $36 \pm 3$  to  $39 \pm 3$  except at 285.1mg/l level of concentration where it exhibited  $40 \pm 2$ .

At 72 and 96 hr. of exposure to both pollutants at selected level of concentration the declining trend right from starting point was observed which was not abrupt rather in a systematic manner except in case of  $CuSO_4$  pollutants at 88mg/level of concentration where it exhibited  $46 \pm 2$ .

The opercular frequency per minute in relation to exposure hr. and different concentration of both pollutant i.e.  $CuSO_4$  and  $K_2Cr_2O_7$  with percent change in case of *C. reba* has been enumerated in Table-2.

In the present study, both fishes when exposed to both heavy metals at different concentration and exposure hour, it was observed that both fishes become sluggish and later on they started erratic swimming losing their equilibrium, loss of balance and sinking to bottom or floating on the surface just prior to death. Almost similar symptoms have been reported by several workers while working on different fishes when treated with different pollutants. Grant and Mehrle (1970) in *C. auratus*, Munsigeri and David (2003) in *C. mrigala*, Prasad et al. (2002) in *C. marulius*, Prasad et al. (2003) in *C. gachua*, S. Thangamalathi et al. (2020) in *C. punctatus* and *Oreochromis niloticus*, Roy et al. (2020) in *C. reba*, Hameed et al. (2005), Gurusamy (2020), Sellers (1975) etc.

It can be safely concluded that the toxicant from the environment mainly enter fish by means of their respiratory system. A mechanism of toxicant uptake through gills probably occurs through pores by simple diffusion and is then absorbed through cell membranes. From the result it is clearly evident that decrease in oxygen consumption by the whole animal may be due to the respiratory distress which affect the opercular movement.

**Conflict of Interest:** The author(s) declares no conflict of Interest.

## REFERENCES

- APHA (1995) standard Methods of the Examination of water and waste water. 14<sup>th</sup> Edn. washington, D.C
- Doudoroff B, Anderson BG, Budick CE, Gattstoff PS, Hart WB, Patrick P, strong ER Surber FW and vanItorn WM (1951) In "Environmental pollution by pesticides" (Ed. Edwards, C.A) plenum press, London and New York, pp. 218
- Grant BF and Mehrle PM (1970) Chronic endrin poisoning in gold fish, *Carassius auratus*. *J. Fish Res. Bd. can* 27: 2225-2231

- Gurusamy K and Ramadas V (2000) Impact of DDT on oxygen consumption and opercular activity of *Lepidocephalichthys thermalis*. *J. Ecotoxicol. Environ. Monit* 10 (4) 197-207.
- Health AG (1987) water pollution and fish physiology CRC, BOCA, Raton FL.USAPP 245.  
<https://doi.org/10.1201/9780203718896>
- Hughes GM, Singh BN (1970) Respiration in an air breathing fish, the climbing perch, *Anabas testudineus* Bloch I. O<sub>2</sub>-uptake and Co<sub>2</sub> release into air water. *J. Exp. Biol.* 94 :265-280.
- Monazaki S et. al. (2015) Xenobiotics and Boremediation (Review), *Nature and science* 13 (2) : 13-115
- Mushigeri SB and Devid M (2003) Assessment of fenvalerate toxicity on oxygen consumption and ammonia excretion in the fresh water fish, *Cirrhinus mrigaia*. *J. Ecotoxicol. Environ, Monit.* 13(3) : 191-195.
- Prasad BB and RoyMK (2003) Effects of copper sulphate and potassium Dichromate on oxygen consumption of the air breathing fish *Channa gachua*. *J. Ecotoxicol. Environ. Monit* 16 (1) 71-76.
- Sellers CM Health, G. and Boss, M.L. 1975: The effects of sublethal concentrations of copper and zinc on ventilation activity, blood oxygen and P<sup>H</sup> in rainbow trout, *salmo gairdenri*. *Water Res.* 9 401-408.
- Singh KM and Prasad, BB (1999) Effects of cadmium chloride and nickel sulphate on *Channa marulius* and *Cirrhinus reba*. *Flora and Fauna*, Vol.5 (1): 13-14.
- Singh SR and Singh BR (1979) Changes in oxygen consumption of a siluroid fish, *Mystus vittatus* put to different concentrations of some heavy metal salts. *Ind. J. Exp. Biol.* 17: 274-246
- Thanga Malathi S and Anuradha V (2020) Lithium induced toxicity profile of oxygen consumption, Haematological parameters and Biochemical profiles of *Channa punctatus* and *Oreochromis niloticus*. *Nature Environment and Pollution Technology* Vol. 19. No. 02 PP. 677-685.
- Vutukuru SS (2000) Chromium induced alterations in some biochemical profiles of the Indian major carp, *Labeo rohita* (Hamilton). *Bull. Environ-contam. Toxicol.* 70, 118-123.